




Antiorbital Ionic Calcium Carbonate (AIC)

Starving Multiple Myeloma by
Re-engineering the Bone Microenvironment

Research by

UAMS
University of Arkansas for Medical Sciences

MOFFITT
CANCER CENTER


Research Funded by

NIH NATIONAL CANCER INSTITUTE

PAULA & RODGER **RINEY**
FOUNDATION


The Multiple Myeloma Paradigm: Seed vs. Soil

The Standard Approach: Attacking the Seed



Marginal effects on myeloma progression, high resistance, inevitable relapse. The soil remains hospitable to surviving cancer cells.


The AIC Innovation: Fortifying the Soil



Ecological modification. Fortifying the bone marrow microenvironment makes it impossible for 5TGM1 myeloma cells to take root or thrive.


Key Insight: Multiple Myeloma heavily relies on the surrounding bone marrow microenvironment. Mice surviving initial transplantation exhibit complete resistance to tumor rechallenge. Fortified bone provides a natural protective mechanism.

Fortifying the Bone Niche: AIC Dietary Intervention to Prevent Multiple Myeloma




The Innovation

Antiorbital Ionic Calcium Carbonate (AIC) is a uniquely engineered, highly bioavailable calcium supplement utilizing weak sigma anti-bonding.



The Mechanism


AIC drives a massive shift in the bone microenvironment, significantly elevating osteoblast differentiation while suppressing osteoclasts.



The Clinical Outcome


Prophylactic AIC treatment radically delays Multiple Myeloma (MM) engraftment and tumor progression in vivo, extending median survival indefinitely in murine models.

Clinical Regimen Design: Pre-Emergence vs. Post-Emergence



Methodology Note: 5TGM1-Luc cells injected via tail vein into NSG (immunodeficient) mice, monitored weekly via bioluminescence.

Survival Outcomes: Complete Prevention of Mortality in NSG Models



PBS Control

- 100% mortality.
- Median survival = 39 days.

AIC TaT


- Delayed mortality.
- Median survival = 45 days ($p = 0.0437$).

AIC TbT

- Unprecedented survival.
- Median survival = Not reached ($p = 0.007$).

THE CONCLUSION: Pre-fortifying the bone environment with AIC before cancer introduction essentially nullifies the lethality of the 5TGM1 MM cells in this model. Even post-emergence intervention showed statistically significant benefit ($p=0.0437$); however, the prophylactic window is clearly the optimal use case.

The Hostile Soil Cross-Section: Shifting the Cellular Balance




Structural Remodeling: The Bone Microenvironment Dashboard

Visual Proof: 3D Micro-Architecture

Control (PBS)




AIC (TbT)




Biochemical Proof: Mineralization

PBS Control



AIC Sustained



Key Metrics (In Vivo Structural Osteogenesis)

Bone Mineral Density (BMD)

Significant Increase
($p=0.030$)

Bone Volume (BV/TV)

Massive Expansion
($p<0.018$)

Trabecular Thickness (Tb.Th)


Structural Fortification
($p<0.022$)

Takeaway: Elevated systemic calcium directly drives massive in vivo and in vitro structural osteogenesis, transforming porous environments into impenetrable mineral matrices.


Disease Progression: AIC Dramatically Delays Tumor Burden

In the NSG immunodeficient mouse model, control subjects exhibited rapid MM engraftment and progression. Conversely, pre-treatment with AIC suppressed tumor growth entirely or delayed it massively.

Bioluminescence Imaging: Tumor Engraftment and Progression



Quantitative Analysis: Total Flux Scatter Plot




Validation: Success in an Immune-Competent Model

To ensure results were not anomalies of immunodeficient mice, the study was replicated in the C57BL/KaLwRij conventional MMBD model—known for pronounced osteolytic lesions and immune competence.


Validation Dashboard

1. Survival




TbT median survival remains **Undefined** vs 46 days for control ($p=0.007$).

2. Bone Volume



Massive, statistically **significant increase** in trabecular bone volume ($p=0.0001$).

3. Cellular Shift



The **exact same biological mechanism** was observed: increased osteoblasts, decreased osteoclasts.

Takeaway: The efficacy of AIC is robust and translatable across diverse biological systems.

Clinical Implications & The Future of Preventative Care



Highly Tolerable Intervention

As a dietary supplement, AIC acts with extreme potency but zero observed renal toxicity or crystallization, unlike high-dose standard calcium.




The Prophylactic Paradigm

Data explicitly proves that pre-treatment (TbT) is vastly superior to post-treatment. Altering the bone niche before cancer emerges is a viable new therapeutic philosophy.



Target Populations

AIC represents a breakthrough, low-toxicity regimen specifically suited for high-risk patients with pre-emerging MM (MGUS/SMM) or for maintaining deep remission and preventing relapse.




THE PARADOX:

In Vitro, AIC has marginal direct effects (<10% inhibition) on isolated myeloma cells. Yet, In Vivo, it stops the cancer entirely.


Why?

The AIC Innovation: Weak Sigma Anti-Bonding

Conventional Calcium




AIC Calcium



- ✓ Independent of Vitamin D for bloodstream absorption.
- ✓ Bypasses Albumin protein binding.
- ✓ Remains as bioavailable ionized calcium.


Antiorbital Ionic Calcium Carbonate (AIC- CaCO_3) fundamentally alters the absorption pathway by altering the molecular bond, maintaining a highly reactive, ionized state in the body. AIC sustains elevated serum ionic calcium levels, significantly restoring calcium homeostasis at both the systemic and cellular levels by stimulating bone formation and cellular decalcification.

Introducing Antiorbital Ionic Calcium Carbonate (AIC)



AIC vs. Conventional Calcium Carbonate (CaCO3)

Dimension	CaCO ₃	AIC
Molecular Bond	Stable Ionic	Sigma Anti-Bonding
Solubility	Low	~200x more soluble
Reaction Rate	0.035 mol/L per sec	0.160 mol/L per sec (4.57x faster)
In Vivo Ionization	Negligible serum change	Rapid, sustained increase in serum Ca²⁺




Long-Term Maintenance

Continuous twice-daily intake yields a sustained elevation of serum calcium ion levels across multiple mouse strains (NSG and C57BL/6)


Key AIC Triggers

- PTH inhibition
- BMP-2 Activation
- Calcitonin Secretion



Elevated urinary calcium is safely excreted. *Ex vivo* ultrasound confirms absolutely zero crystallization or kidney stone formation after prolonged AIC regimens.

The Molecular Blueprint: Sigma Anti-Bonding



The anti-bonding orbital configuration creates exceptional water molecule solvation-driven coordination, allowing AIC to instantly flood the system with active calcium ions.



Oncology: Normalizing the Tumor Microenvironment

AIC does not act as indiscriminate cytotoxic chemotherapy; it alters the biochemical “soil” so the cancer “seed” cannot thrive.


Blockade 1: Shutting Down Survival Signals

(IL-6 / TNF- α / IL-1 β)

Prevents Tumor-Associated Macrophages (TAMs) from turning on anti-apoptosis switches that keep cancer cells alive.

Kyung Hee University's AIC research on extracellular matrix (ECM) protection and inflammation reduction effectively complements our MM study.

TME Spatial Diagram



Blockade 3: Halting Metastatic Escape

(MMPs)

Strengthens structural tissue barriers, disabling the molecular scissors cancer uses to enter the bloodstream.

Blockade 2: Blocking Angiogenesis Fuel Lines

(COX-2)

Prevents the tumor from signaling the body to build dedicated blood vessels for nutrient supply.

Human Case Study #1: Initiation of AIC Following Discontinuation of Conventional Treatments



Patient 01

Female, Age 80, South Korea

- Diagnosed with Multiple Myeloma in 2012.
- Commenced AIC Therapy in Dec. 2013.

Cancer Remission (Hematology)

Baseline (Nov 2013)
WBC 2,190 /uL



Current (Jan 2015)
WBC 4,760 /uL

Platelets and Neutrophils fully normalized.

Renal Recovery

Baseline (Nov 2013)
eGFR 12 ml/min



Current (Nov 2015)
eGFR 62 ml/min

Reversal of renal impairment

Bone Density (Osteoporosis Reversal)

Baseline (Nov 2013)
T-Score -3.6



Current (Nov 2015)
T-Score -0.9

(Normal Range: Above -1.0).

This clinical case was provided by ACRI. The patient achieved a normal CBC within 14 months of initiating AIC therapy. Furthermore, her previously declining kidney function normalized, and her osteoporosis was fully reversed.

Human Case Study #2: Initiation of AIC for 5 months Prior to Conventional Treatments



Patient 02
Female, Age 67,
USA

- Diagnosed Nov. 2023. Presented with diffuse lytic lesions and multiple compression fractures. Refused to see an oncologist.
- No conventional MM treatments received for 5 months, including Zometa. Maintained exclusively on AIC Therapy (3-4x daily) and pain medication.

Hypercalcemia Resolution

Baseline (Nov 2023):
Calcium 11.2 mg/dL

DOWN

Current (Apr 2024):
Calcium 9.1 mg/dL

Return to Normal Range (8.4-10.5).

Renal Function Restoration

Baseline (Mar 2024):
E-GFR 62 mL/min

UP

Current (Apr 2024):
E-GFR 83 mL/min

Strong function (Normal: ≥ 60).
Total Urine Protein dropping sharply (24 to 9.6).

Systemic Resilience

Recovery trajectory maintained despite severe intervening infections between Jan-Feb 2024 (Meningitis, Pneumonia, C. difficile).




This clinical case was provided by ACRI. The patient began conventional treatments (Velcade and Darzalex, without Zometa) in May 2024, following improvement with AIC therapy. As of 2026, on her oncologist's recommendation, she has discontinued all treatments except for a daily maintenance dose of AIC therapy. She continues to be actively monitored.

The Synthesis:



The AIC Microenvironment Framework



Chemistry: Sigma Anti-Bonding Configuration creates a massive influx of bioavailable Ca^{2+} ions.



Outcome: Tumor starvation, failed engraftment, and undefined survival extension.



Physiology: Ion flood triggers an Osteoclast-to-Osteoblast cellular shift, radically increasing trabecular density.



Oncology: The remodeled, structurally dense bone niche becomes fundamentally **hostile** to multiple myeloma colonization.



To book a Zoom session with ACRI research team to learn more about AIC therapy.



To study more research and clinical cases of AIC therapy.



ADVANCED
CALCIUM
RESEARCH
INSTITUTE

www.aictherapy.com